

EGS Abstract for The Hague, 1999

SUBSTORM PSEUDOBREAKUPS ASSOCIATED WITH INTER-PLANETARY SHOCKS/PRESSURE PULSES: WIND AND POLAR

X.-Y. Zhou (1), B. T. Tsurutani (1), J. K. Arballo (1), D. Berdichevsky (2) and R. Lepping (2)

(1) Jet Propulsion Laboratory, California Institute of Technology, Pasadena, CA 91109, USA, (2) Goddard Space Flight Center, Code 695.0, Greenbelt, MD 20771, USA.

zhou@jpl.nasa.gov/Fax: 818 354-8895

Ten interplanetary WIND shock/ pressure pulse events are used to study the magnetospheric delay of pseudobreakup (PB) or substorm onsets. We identify the PBs and substorms by using the **POLAR UV imaging data**. The states of the interplanetary medium and the conditions of the ionosphere before and after the **auroral brightening** onsets are studied. We find that the magnetospheric delay time strongly constrains the location of the nightside X-line during such events. We also find for PB (or no activity) events, that the interplanetary and ionosphere preconditions are unusually low.

Submittal Information

- | | |
|---------------------------|--|
| 1. Zhou | 2. Session ST9-02, Auroral and inner magnetosphere |
| 4800 Oak Grove Drive | 3. M. Grande and H.J. Opgenoorth |
| Tel.: 818 354-9169 | 4. none |
| Fax.: 818 354-8895 | 5. Oral presentation strongly preferred |
| E-mail: zhou@jpl.nasa.gov | |

Abstracts to be submitted on or before December 15, 1997 to

EGS Office
Max-Planck-Str. 13
37191 Katlenburg-Lindau
Germany

Tel.: [+49] 5556-1440
Fax.: [+49] 5556-4709
Email: EGS@Copernicus.org
<http://www.copernicus.org/EGS/EGS.html>